

What we need from XG observatories to unveil the birth, life and death of massive stars – Part 3: stellar death

SUMMARY

The next generation of gravitational-wave (GW) observatories have the potential to solve key questions about massive stellar evolution. In this last part, we discuss how next-generation measurements of the remnant mass distribution can constrain supernova physics and nuclear reaction rates in the final stages of a massive star's life. To achieve this we need to push our detection frequency down to 5 Hz and detect 10,000 double compact object (DCO) mergers with masses below $10M_{\odot}$ and a S/N above 100.

Key questions in massive binary stellar evolution: 3. How do massive stars end their lives?

Massive stars impact *every* part of modern astrophysics; their ejecta, shocks, outflows, and ionizing photons shape their environments, they trigger and regulate star formation, and drive the chemical evolution of the Universe that enables the formation of elements like oxygen, and the more complex molecules necessary to facilitate life. Despite their importance, the formation, lives, and explosive deaths of massive stars is still a mystery. They are rare and short-lived, making it extremely challenging to observe a statistically significant population and learn about their properties, especially in environments outside our Milky Way. GW astrophysics provides a new frontier to study the lives and deaths of massive stars throughout cosmic history and can help solve key questions in massive star evolution: 1. *How do massive stars form?*, 2. *How do massive stars evolve and interact?*, and 3. *How do massive stars end their lives?*.

How do massive stars end their lives? Modeling the final moments of a massive star's life presents a significant challenge, yet it is crucial for our understanding of phenomena such as supernovae (SN), long-duration gamma-ray bursts, and pulsational pair-instability supernovae (PPISN) that are prevalent throughout our Universe. Extensive research has been conducted on SN explosions, but the precise mechanism behind stellar core collapse and the physical engine driving the subsequent explosion remain elusive. Moreover, it is uncertain whether BHs experience significant natal kicks and to what extent material falls back when a star explodes, leading to the formation of a neutron star (NS) or a black hole (BH).

Potential scientific impact of XG detectors

Accurate measurements of the mass distribution of NS and BHs offer invaluable insights into the aforementioned topics. Next-generation detectors present a unique opportunity for significant advancements in several ways:

Firstly, Xg ground-based detectors will provide access to much lower frequencies, reaching as low as 5Hz compared to the current limit of 20Hz for present-day detectors [14, 3, 4]. This allows us to observe more cycles of the inspiral phase of massive events (such as for GW190521 [1]), removing doubt about their observed source properties. Moreover, this will enable us to detect BBH merger with masses up to $M_{tot} \leq 450M_{\odot}$ (instead of $M_{tot} \leq 110M_{\odot}$). Mapping the mass distribution of merging BBHs with masses of $M_{tot} \approx 100 - 400M_{\odot}$ in detail will reveal *both edges* of the PISN mass gap, which can be used to measure the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ reaction rate [5]. This measurement, in turn, can be used to refine our predictions for the lower end of the mass spectrum.

Secondly, there will be no bias towards higher-mass systems since typical sources will not be detected at threshold [19]. This will allow us to place unprecedented constraints on the source properties of NSNS, NSBH, and low-mass BHBHs. Currently, due to low signal-to-noise (S/N) observations, our studies are restricted to analyzing chirp mass or total mass. However, the anticipated increase in the number of extreme S/N detections facilitated by next-generation detectors will allow us to precisely determine the mass ratio and, consequently, the component masses [3]. This will facilitate a detailed measurement of the mass distribution of low-mass double compact objects (DCOs), which will allow us to distinguish between various proposed theories that seek to explain the shape of the remnant mass distribution, and with it the long-debated existence of a NS-BH mass gap [2, 8, 15, 6, 13, 20, 10, 17]. These theories include the fallback mechanism [7, 9], the failed supernova scenario [12, 11], binary evolutionary effects [18], or details in the compactness of stellar cores at core collapse [16].

SCIENTIFIC IMPACT OF XG DETECTORS

The first measurement of both edges of the PISN mass gap, and a corresponding measurement of the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ nuclear reaction rate. Constraints on the abundances of intermediate-mass BHs. Detailed (sub solar mass precision) measurement of the low-mass distribution of NSNS, and BHBH component masses, and consequently new constraints on the physics core-collapse SNe.

Benchmarks for XG detectors to enable the scientific impact

- 1. A minimum detection frequency that reaches down to 5Hz, to constrain the mass distribution above the PISN-mass gap.** To constrain the mass distribution above the PISN-mass gap, it is crucial to achieve a minimum detection frequency of 5 Hz. By attaining this level of sensitivity, we can uncover the width and location of the PISN gap down to 5% accuracy, which will, in turn, allow a measurement of the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ nuclear reaction rate. Additionally, lower frequencies enable us to observe more cycles of all higher-mass BHBH mergers, thereby reducing source property uncertainties associated with high-mass systems. Moreover, this will allow us to approach the population of intermediate-mass BHs from the stellar-mass side, shedding the first light on this previously unknown population.
- 2. 10,000 detections with $\text{S/N} > 100$ and masses below $10M_{\odot}$, to probe supernova physics.** Systems with low masses and/or extreme mass ratios are particularly disfavored by current-day detector networks. Yet they will be crucial to distinguish between different proposed scenarios for the shape of the remnant-mass distribution and the corresponding supernova physics. We need a large set of systems with high SNR events with low masses and/or high mass ratios, such that we can place exact constraints on the component masses. This is achievable with XG detectors [14]. Specifically, about 10,000 events with component masses between $1-10M_{\odot}$, would roughly lead to a precision of 3% per bin of 1000 events.

XG DETECTOR AND NETWORK REQUIREMENTS

A network of detectors that extends to lower frequencies of about 5Hz, and with a noise floor that is a factor 10 lower than the capabilities of current detectors.

Authors

L. A. C. van Son^{1,2}, lieke.van.son@cfa.harvard.edu

F. S. Broekgaarden², floor.broekgaarden@cfa.harvard.edu

¹Center for Astrophysics | Harvard & Smithsonian, 60 Garden St., Cambridge, MA 02138, USA,

²Anton Pannekoek Institute for Astronomy, University of Amsterdam, Science Park 904, 1098XH Amsterdam, NL

Bibliography

- [1] R. Abbott et al. “Properties and Astrophysical Implications of the 150 M_{\odot} Binary Black Hole Merger GW190521”. In: *ApJ* 900.1, L13 (Sept. 2020), p. L13. DOI: [10.3847/2041-8213/aba493](https://doi.org/10.3847/2041-8213/aba493). arXiv: [2009.01190](https://arxiv.org/abs/2009.01190) [[astro-ph.HE](#)].
- [2] Charles D. Bailyn et al. “The Mass Distribution of Stellar Black Holes”. In: *ApJ* 499.1 (May 1998), pp. 367–374. DOI: [10.1086/305614](https://doi.org/10.1086/305614). arXiv: [astro-ph/9708032](https://arxiv.org/abs/astro-ph/9708032) [[astro-ph](#)].
- [3] Marica Branchesi et al. “Science with the Einstein Telescope: a comparison of different designs”. In: *arXiv e-prints*, arXiv:2303.15923 (Mar. 2023), arXiv:2303.15923. DOI: [10.48550/arXiv.2303.15923](https://doi.org/10.48550/arXiv.2303.15923). arXiv: [2303.15923](https://arxiv.org/abs/2303.15923) [[gr-qc](#)].
- [4] Matthew Evans et al. “A Horizon Study for Cosmic Explorer: Science, Observatories, and Community”. In: *arXiv e-prints*, arXiv:2109.09882 (Sept. 2021), arXiv:2109.09882. DOI: [10.48550/arXiv.2109.09882](https://doi.org/10.48550/arXiv.2109.09882). arXiv: [2109.09882](https://arxiv.org/abs/2109.09882) [[astro-ph.IM](#)].
- [5] R. Farmer et al. “Constraints from Gravitational-wave Detections of Binary Black Hole Mergers on the $^{12}\text{C}(\alpha, \gamma)^{16}\text{O}$ Rate”. In: *ApJ* 902.2, L36 (Oct. 2020), p. L36. DOI: [10.3847/2041-8213/abbadd](https://doi.org/10.3847/2041-8213/abbadd). arXiv: [2006.06678](https://arxiv.org/abs/2006.06678) [[astro-ph.HE](#)].
- [6] W. M. Farr et al. “The Mass Distribution of Stellar-mass Black Holes”. In: *ApJ* 741, 103 (Nov. 2011), p. 103. DOI: [10.1088/0004-637X/741/2/103](https://doi.org/10.1088/0004-637X/741/2/103). arXiv: [1011.1459](https://arxiv.org/abs/1011.1459).
- [7] C. L. Fryer et al. “Compact Remnant Mass Function: Dependence on the Explosion Mechanism and Metallicity”. In: *ApJ* 749, 91 (Apr. 2012), p. 91. DOI: [10.1088/0004-637X/749/1/91](https://doi.org/10.1088/0004-637X/749/1/91). arXiv: [1110.1726](https://arxiv.org/abs/1110.1726) [[astro-ph.SR](#)].
- [8] Chris L. Fryer and Vassiliki Kalogera. “Theoretical Black Hole Mass Distributions”. In: *ApJ* 554.1 (June 2001), pp. 548–560. DOI: [10.1086/321359](https://doi.org/10.1086/321359). arXiv: [astro-ph/9911312](https://arxiv.org/abs/astro-ph/9911312) [[astro-ph](#)].
- [9] Chris L. Fryer, Aleksandra Olejak, and Krzysztof Belczynski. “The Effect of Supernova Convection On Neutron Star and Black Hole Masses”. In: *ApJ* 931.2, 94 (June 2022), p. 94. DOI: [10.3847/1538-4357/ac6ac9](https://doi.org/10.3847/1538-4357/ac6ac9). arXiv: [2204.13025](https://arxiv.org/abs/2204.13025) [[astro-ph.HE](#)].

- [10] Peter G. Jonker et al. “The Observed Mass Distribution of Galactic Black Hole LMXBs Is Biased against Massive Black Holes”. In: *ApJ* 921.2, 131 (Nov. 2021), p. 131. doi: [10.3847/1538-4357/ac2839](https://doi.org/10.3847/1538-4357/ac2839). arXiv: [2104.03596](https://arxiv.org/abs/2104.03596) [[astro-ph.HE](#)].
- [11] C. S. Kochanek. “Constraints on core collapse from the black hole mass function”. In: *MNRAS* 446.2 (Jan. 2015), pp. 1213–1222. doi: [10.1093/mnras/stu2056](https://doi.org/10.1093/mnras/stu2056). arXiv: [1407.5622](https://arxiv.org/abs/1407.5622) [[astro-ph.SR](#)].
- [12] C. S. Kochanek. “Failed Supernovae Explain the Compact Remnant Mass Function”. In: *ApJ* 785.1, 28 (Apr. 2014), p. 28. doi: [10.1088/0004-637X/785/1/28](https://doi.org/10.1088/0004-637X/785/1/28). arXiv: [1308.0013](https://arxiv.org/abs/1308.0013) [[astro-ph.HE](#)].
- [13] Laura Kreidberg et al. “Mass Measurements of Black Holes in X-Ray Transients: Is There a Mass Gap?” In: *ApJ* 757.1, 36 (Sept. 2012), p. 36. doi: [10.1088/0004-637X/757/1/36](https://doi.org/10.1088/0004-637X/757/1/36). arXiv: [1205.1805](https://arxiv.org/abs/1205.1805) [[astro-ph.HE](#)].
- [14] Michele Maggiore et al. “Science case for the Einstein telescope”. In: *JCAP* 2020.3, 050 (Mar. 2020), p. 050. doi: [10.1088/1475-7516/2020/03/050](https://doi.org/10.1088/1475-7516/2020/03/050). arXiv: [1912.02622](https://arxiv.org/abs/1912.02622) [[astro-ph.CO](#)].
- [15] F. Özel et al. “The Black Hole Mass Distribution in the Galaxy”. In: *ApJ* 725 (Dec. 2010), pp. 1918–1927. doi: [10.1088/0004-637X/725/2/1918](https://doi.org/10.1088/0004-637X/725/2/1918). arXiv: [1006.2834](https://arxiv.org/abs/1006.2834).
- [16] Fabian R. N. Schneider, Philipp Podsiadlowski, and Eva Laplace. “Bimodal black-hole mass distribution and chirp masses of binary black-hole mergers”. In: *arXiv e-prints*, arXiv:2305.02380 (May 2023), arXiv:2305.02380. doi: [10.48550/arXiv.2305.02380](https://doi.org/10.48550/arXiv.2305.02380). arXiv: [2305.02380](https://arxiv.org/abs/2305.02380) [[astro-ph.HE](#)].
- [17] Jared C. Siegel et al. “Investigating the Lower Mass Gap with Low Mass X-ray Binary Population Synthesis”. In: *arXiv e-prints*, arXiv:2209.06844 (Sept. 2022), arXiv:2209.06844. arXiv: [2209.06844](https://arxiv.org/abs/2209.06844) [[astro-ph.HE](#)].
- [18] L. A. C. van Son et al. “No Peaks without Valleys: The Stable Mass Transfer Channel for Gravitational-wave Sources in Light of the Neutron Star-Black Hole Mass Gap”. In: *ApJ* 940.2, 184 (Dec. 2022), p. 184. doi: [10.3847/1538-4357/ac9b0a](https://doi.org/10.3847/1538-4357/ac9b0a). arXiv: [2209.13609](https://arxiv.org/abs/2209.13609) [[astro-ph.HE](#)].
- [19] Salvatore Vitale. “Three observational differences for binary black holes detections with second- and third-generation gravitational-wave detectors”. In: *Phys. Rev. D* 94.12, 121501 (Dec. 2016), p. 121501. doi: [10.1103/PhysRevD.94.121501](https://doi.org/10.1103/PhysRevD.94.121501). arXiv: [1610.06914](https://arxiv.org/abs/1610.06914) [[gr-qc](#)].

- [20] Lukasz Wyrzykowski and Ilya Mandel. “Constraining the masses of microlensing black holes and the mass gap with Gaia DR2”. In: *A&A* 636, A20 (Apr. 2020), A20. doi: [10.1051/0004-6361/201935842](https://doi.org/10.1051/0004-6361/201935842). arXiv: [1904.07789](https://arxiv.org/abs/1904.07789) [[astro-ph](https://arxiv.org/archive/astro).SR].